

WHAT ARE THE HEALTH FACILITY FACTORS INFLUENCING MALE INVOLVEMENT IN ANTENATAL CARE AMONG CLIENTS ATTENDING ANTENATAL CLINIC AT NEBBI GENERAL HOSPITAL, NEBBI DISTRICT?

Teddy Adero^{*1}, Atim Miria²

Department of Nursing and Midwifery, Jerusalem Institute of Health Sciences

ABSTRACT

Background

Antenatal care (ANC) is crucial for maternal and fetal health, yet male involvement in ANC remains low globally. Despite efforts to encourage male participation, studies indicate low levels of involvement across Africa, including Uganda. This study aims to identify factors influencing male involvement in ANC at Nebbi General Hospital.

Methods

The study employed a cross-sectional quantitative approach using purposive sampling to collect data from pregnant women and their partners attending the antenatal clinic, with a final sample size of 37 participants, and data analysis was conducted using SPSS version 20.0

Results and discussions

There was 100% response rate with 37 participants, revealing distance to the facility, discussion of antenatal care with health workers, and comfort with health workers' gender as key determinants to male involvement.

Conclusion

Various health facility factors significantly influence male participation in antenatal care (ANC), affecting maternal and child health outcomes.

Recommendation

To improve male involvement in ANC, hospitals should provide flexible clinic hours, especially for working men. District health officers should address health facility factors and train health workers to engage with male partners effectively.

Nursing implications

Nurses should educate expectant mothers and their male partners on the importance of male involvement in ANC and its impact on pregnancy outcomes, advocating for flexible clinic hours to accommodate diverse backgrounds.

KEY WORDS: Antenatal care, Male involvement, Nebbi General Hospital.

Submitted: January 01, 2025 Accepted: March 20, 2025 Published: June 01, 2025

Corresponding author: Adero Teddy

Email: tyadero@gmail.com

Department of Nursing and Midwifery, Jerusalem Institute of Health Sciences

BACKGROUND

Antenatal care (ANC) is critical for the health and development of the fetus because it establishes a connection between the mother and her family and the healthcare

system, potentially increasing the likelihood that a skilled attendant will be used at birth and promoting good health throughout the life cycle (Annie, 2021). ANC from a qualified health provider is essential to screen pregnancy and decrease the danger of morbidity for mother and baby

during pregnancy, delivery, and the postnatal period (WHO, 2015). The concept of male involvement in maternal well-being is currently being promoted as a fundamental component of World Health Organization (WHO) efforts to make pregnancy safer (Manda-Taylor et al., 2017). Traditionally, ANC has been perceived as a responsibility primarily for women, with male partners often marginalized from the process. However, there is an increasing recognition of the importance of involving men in ANC to enhance maternal and child health outcomes (Kirui, 2021).

Like in most other African nations, family planning, getting pregnant, and giving birth have traditionally been seen as matters entirely about women in South Africa (Annie, 2021). However, social and sexual domination by males can seriously increase a woman's chance of infection and unintended pregnancy. Additionally, throughout a pregnancy, a man's sexual behavior can have an impact on the health of both the mother and the unborn child (Kirui, 2021).

Globally, men continue to have a low level of involvement in mother and child health in both developed and developing nations, though the World Health Organization advocates for male participation in ANC as a crucial component of a comprehensive approach to maternal and child healthcare (Nyasiro S et al, 2019).

Similarly, in Africa, a similar trend of low male involvement is observed; for instance, in Ethiopia, 25% of expectant mothers have their husbands accompany them for ANC (Mamo et al., 2017). In Tanzania, it was also observed that there is a low proportion of male engagement of 56.9% (Kabanga et al., 2019). In addition, reports from Kenya show that male involvement is at 34.1% (Muia et al., 2022).

According to MoH (2020), maternal mortality ratio in Uganda remains high with 336 maternal deaths per 100,000 live births, Infant mortality is 43 deaths per 1000 live births, with 42% of the mortality occurring during the neonatal period of which most of them is as a result of low involvement of males who could assist in many ways (Babughirana, 2020). However, there is an increasing number of initiatives and programmatic efforts to come up with strategies that encourage male involvement in safe motherhood (Yargagwa & Leonardi-Bee, 2015). In Malawi and Uganda, women who were accompanied by their spouses were given priority in service provision as a strategy to encourage and support male participation in utilization of ANC services (Atuahene et al., 2017). This trend of low male involvement, according to Byamugisha, et al. (2017), has been attributed to different factors which have been identified in other studies as: Health-facility

factors, Cultural factors, and Socio-Economic factors. According to Auma et al. (2023), a study was carried out in the Palabek Refugee Settlement, Lamwo district, Northern Uganda, to evaluate the factors associated with male involvement in ANC. The prevalence of male participation in ANC was 39%. Anecdotal information from Nebbi District Hospital indicates a low level of male involvement in antenatal care services, with only 9 males attending ANC services at Nebbi General Hospital during the financial year 2022-2023. However, there is generally limited information to explain the poor involvement of males in ANC services. Therefore, the study aims to identify factors influencing male involvement in antenatal care among clients attending the antenatal clinic at Nebbi General Hospital, Nebbi district.

METHODOLOGY

Introduction

This chapter focuses on the study design and rationale, study setting and rationale, study population, sample size determination, sampling procedure, inclusion criteria, definitions of variables, research instrument, data collection procedures, data management, data analysis, ethical considerations, limitations of the study, and dissemination of results.

Study design and rationale

This study was a cross-sectional study using a quantitative approach to determine factors influencing male involvement in antenatal care among clients attending the antenatal clinic at Nebbi General Hospital, Nebbi district. The selected design was optimal for data collection at a specific moment in time, as the data were collected at a specific point.

Study setting and rationale

The study was conducted among clients attending the antenatal clinic at Nebbi General Hospital, Nebbi district. The hospital offers many health care services, including immunization, child health services, HIV/AIDS management services, general patient management, surgery, laboratory services, nutrition services, antenatal, maternity, and post-natal services, EMTCT program, as well as RCT services, among many others.

The hospital is located in the central business district of the town of Nebbi, in Nebbi District, in the West Nile sub-region, in Northern Uganda, about 78 kilometers (48 mi) southeast of Arua Regional Referral Hospital. The ANC unit is usually run by 5 staff, and it operates from Monday to Friday with a total of 4 examination beds used for the examination of the mothers. The study setting is chosen because it is easily accessible to the researcher and has a specific population.

Study population

The study was conducted among pregnant women and their partners attending the antenatal clinic at Nebbi General Hospital, Nebbi district, at the time of the study.

Sample size determination

The sample size was determined using the single-proportion formula of Fischer's et al as follows:

$$n = (Z \alpha/2)^2 P (1-P)/e^2$$

Where;

n = sample size needed

Z $\alpha/2$ =level of statistical significance at 95% confidence interval (standard value 1.96)

P= proportion of males involved in ANC services, i.e., 10%
e=maximum acceptable marginal error- 5% (0.05)

$$\text{Therefore; } n_0 = 1.962 (0.1) (1-0.1) / (0.05)^2 = 138.3$$

$$n_0 = 139 \text{ participants}$$

Using the finite population factor for sample size adjustment by Glenn D. Israel 1994

$$n = \frac{n_0 \times N}{n_0 + (N-1)}$$

Where: N = total number of clients attending ANC in a day
=50

$$n = \frac{139 \times 50}{139 + (50-1)} = 36.96$$

$$n = 37 \text{ participants}$$

Therefore, the study participants were 37. This is above the minimum number expected by UNMEB and is representative of the entire population.

Sampling procedure

Purposive sampling is a non-probability sampling technique where researchers intentionally select participants or cases available at the facility at the time of study. This was chosen since Nebbi Hospital is a high-volume facility, which serves clients from both rural and urban settings.

Inclusion criteria

All the pregnant women and their partners attending the antenatal clinic at Nebbi general hospital, Nebbi district, at the time of study, who consented to take part.

Exclusion criteria

All patients who could not participate in the study, insane clients, those who refused to consent to the study, and clients under 18years.

Definition of Variables

Dependent variable

The dependent variable is the outcome or response variable that is being observed or measured.

The dependent variable was male involvement in Antenatal Care.

Independent variable

An independent variable is a variable that is manipulated or controlled by the researcher.

The independent variable was the factors influencing, i.e., health facility factors, etc.

Research instrument

Self-administered structured questionnaires were distributed to gather data on the factors linked to male involvement among clients attending ANC at Nebbi General Hospital, Nebbi district. The questionnaire was composed of four parts, i.e., part 1- introduction, part 2 health facility factors. A pilot study was conducted on 5 respondents in a single day at Jupangira H/C II to ensure the reliability and validity of the tool. Errors and ambiguous questions were identified and corrected with the help of the supervisor.

Data collection procedure

Following approval, an introductory letter was obtained from the principal of the Jerusalem School of Nursing and Midwifery. The letter was then taken to the Medical Superintendent of Nebbi General Hospital. Subsequently, he introduced the researcher to the ANC ward in charge, who, in turn, introduced the researcher to the respondents. The purpose of the research was clearly explained to the respondents before they consented and filled out the questionnaires. Data collection spanned approximately four (4) days, during which a total of 10 respondents were obtained per day.

Data management

All study participants were given a unique identification number that was recorded in the questionnaire. The hard copies of the interview transcripts were kept under lock and key, and the data were accessible only to the researcher. The questionnaires were carefully examined to verify the precision and thoroughness of the gathered information. Following that, the data were input, encoded, and refined using Statistical Package for Social Sciences (SPSS) version 20.0.

Data analysis

Statistical Package for Social Sciences (SPSS) version 20 was used to conduct the analysis. Descriptive statistics were used in univariate analysis to describe independent variables, which were displayed as frequencies and

percentages, such as age, education level, marital status, occupation, parity, and address. Through cross-tabulation, bivariate analysis will investigate relationships between dependent and independent variables. Testing was done on categorical data (binary, ordinal, or nominal) using Fischer's exact test or Chi-Square (X²). The following numerical (continuous) variables—mode, mean, range, variance, and standard deviation—were evaluated for indicators of central tendencies. Text, tables, and graphs were used to present the findings.

Ethical consideration

Approval

Approval for the study was obtained from the Research and Ethics Committee. Furthermore, authorization was requested from the administrators, i.e., MS Nebbi General Hospital. To formalize this procedure, an official letter issued by the institution was provided to the MS Nebbi General Hospital before the initiation of data collection.

Consent

A written informed consent form, delineating the research's objectives, potential advantages, associated risks, and the rights of the participants, was verbally explained to all respondents. Subsequently, participants were requested to express their consent after confirming their comprehension and willingness to participate in the study. Consent was indicated through a written signature or a thumbprint, particularly for individuals who faced challenges in writing.

Privacy

To ensure privacy, interviews with respondents were conducted in a confidential and secure environment, inaccessible to others. Additionally, all information and data collected were entered into SPSS and promptly secured with a password, enhancing confidentiality and safeguarding the privacy of the participants.

RESULTS

Introduction

This chapter describes the results, which include; health facility factors influencing male involvement in antenatal care among clients attending antenatal clinic at Nebbi general hospital, Nebbi district. A response rate of 100 % was obtained with a total enrolment of 37.

Health facility factors of the respondents

Confidentiality

The data collection was carried out by the researcher personally, and the gathered information was stored in a location with restricted access. Participant confidentiality was upheld by using initials instead of full names in all records. Access to the collected data was restricted to the research team alone, ensuring the confidentiality of participant information.

Limitations of the study


There was information bias arising from the sensitivity of the information, as participants opted to withhold certain details. To address this concern, counseling was implemented to build rapport and encourage openness among participants.

Additionally, financial constraints posed challenges to the research process, given the associated expenses such as transportation to the study site and the purchase of necessary supplies. These limitations had the potential to disrupt the smooth execution of the research.

Dissemination of results

The findings of the study were compiled, and each copy was sent to various recipients. Specifically, copies were forwarded to the Uganda Nurses and Midwives Examinations Board. Another copy was provided to the Jerusalem school of nursing and midwifery to contribute to the enhancement of the school library. The administration of Nebbi General Hospital received a copy to facilitate actions in accordance with the study's recommendations. Additionally, the researcher retained a copy for personal ownership and future reference.

Figure 1 showing distance of facility from the respondents



When considering the distance to the facility, 40.5% (15) of respondents lived within a medium distance, followed by 37.8% (14) who lived nearby, and 21.6% (8) who lived farther away.

Table 2 showing whether distance to the facility is a challenge to the respondent



A significant proportion of respondents (40.5%, 15) found the distance to be a challenge, while the majority (59.5%, 22) did not perceive it as a challenge.

Table 1 showing residence of the respondents

	Frequency	Percent	Valid Percent	Cumulative Percent
Urban	21	56.8	56.8	56.8
Rural	16	43.2	43.2	100.0
Total	37	100.0	100.0	


The majority of respondents (56.8%, 21) lived in urban areas, while 43.2% (16) lived in rural areas.

Table 2 showing respondents' satisfaction with services at the hospital

	Frequency	Percent	Valid Percent	Cumulative Percent
Yes	28	75.7	75.7	75.7
No	9	24.3	24.3	100.0
Total	37	100.0	100.0	

The majority of respondents (75.7%, 28) were satisfied with the services delivered, while a minority (24.3%, 9) were not.

Figure 3 showing whether the respondents discussed ANC findings with the health worker




The majority of respondents (62.2%, 23) had discussed ANC with a health worker, while a minority (37.8%, 14) had not.

Table 3 showing adequacy of space in the facility

	Frequency	Percent	Valid Percent	Cumulative Percent
Yes	32	86.5	86.5	86.5
No	5	13.5	13.5	100.0
Total	37	100.0	100.0	

The majority of respondents (86.5%, 32) found the space adequate, while a minority (13.5%, 5) did not.

Figure 4 showing gender attending to the respondents



Male attended majority that is 91.9% (34) of respondents to by female, while 8.1% (3), by males.

Table SEQ Table | * ARABIC 11 showing comfort with the gender attending to the respondent

Freq	Perc	Vali	Cu
uencent	d	mul	
y	Percativ		
ent	e		
	Perc		
	ent		
Yes	30	81.1	81.1
No	7	18.9	18.9
		100.0	100.0

Tota	37	100.100.
l	0	0

The majority of respondents (81.1%, 30) were comfortable with the gender of the health worker attending to them, while a minority (18.9%, 7) were not.

Table SEQ Table | * ARABIC 12 showing satisfaction with the attitude of health workers by the respondents

Freq	Perc	Vali	Cu
uencent	d	mul	
y	Percativ		
	ent e		
	Perc		
	ent		

Yes	28	75.7	75.7	75.7
-----	----	------	------	------

No	9	24.3	24.3	100.
				0

Tota	37	100.100.
l	0	0

The majority of respondents (75.7%, 28) were satisfied with the attitude of health workers, while a minority (24.3%, 9) were not.

Table SEQ Table | * ARABIC 13 showing access of information at ANC clinic by the respondents

Freq	Perc	Vali	Cu
uencent	d	mul	
y	Percativ		
	ent e		
	Perc		
	ent		

Vali	Yes	24	64.9	64.9	64.9
d					

No	13	35.1	35.1	100.
				0

Tota37 100.100.
1 0 0

The majority of respondents (64.9%, 24) had access to information at the ANC clinic, while a minority (35.1%, 13) did not.

DISCUSSION

Introduction

This chapter presents the discussion of findings, conclusions, and recommendations of the study, which were obtained after data analysis and implications for nursing practice.

Health facility factors influencing male involvement in antenatal care

The majority of respondents (86.5%, 32) find the space adequate, while a minority (13.5%, 5) do not. In the Democratic Republic of the Congo, a study revealed that male involvement in ANC clinics is adversely affected by the insufficient space designated for male partners. The study conducted by Deitkemena et al. (2012) highlighted that clinics often face limitations in accommodating both pregnant patients and their male partners simultaneously due to space constraints. This lack of adequate space not only hinders male partners from actively participating in antenatal care but also underscores the need for services specifically tailored to address men's concerns. Creating a welcoming environment for male partners in ANC clinics, with sufficient space and tailored services, is crucial for promoting their active involvement in maternal health care. The majority of respondents (81.1%, 30) are comfortable with the gender of the health worker attending to them, while a minority (18.9%, 7) are not. Men may feel uncomfortable in health clinics where the majority of the staff and patients are women. This imbalance in gender representation may contribute to a sense of unease or unfamiliarity for men in such settings. Therefore, ensuring gender diversity among health workers and providing education on relevant topics can help create a more welcoming environment for male partners in antenatal care clinics, ultimately promoting their active involvement in maternal health care.

The majority of respondents (62.2%, 23) have discussed Antenatal Care (ANC) with a health worker, while a minority (37.8%, 14) have not. In Tanzania, only 23.5% of men accompanying their partners reported having the opportunity to discuss maternal health issues with healthcare providers, according to the study conducted by Gibore et al. (2019). This suggests a significant gap in engaging male partners in ANC discussions. Increased access to ANC information for men can lead to greater involvement in maternal health care, ultimately

contributing to improved maternal and child health outcomes.

CONCLUSION

Health facility factors such as space adequacy, health worker gender, and ANC discussion opportunities influence male involvement. Understanding and addressing these factors are crucial for promoting male participation in ANC and improving maternal and child health outcomes.

RECOMMENDATION

The hospital should address work-related barriers by providing flexible ANC clinic hours to accommodate men's work schedules, especially for those who work long hours, and offer weekend or evening ANC clinics to make it easier for working men to participate.

Ensuring gender diversity among health workers and providing education on relevant topics can help create a more welcoming environment for male partners in antenatal care clinics, ultimately promoting their active involvement in maternal health care.

The district health officer should improve health facility factors, such as ensuring that ANC clinics have adequate space to accommodate male partners comfortably, train health workers to create a welcoming environment for male partners, and provide them with relevant information and support. It should also encourage discussions about ANC between health workers and male partners to increase their understanding and involvement in maternal health care.

ACKNOWLEDGEMENT

I express my foremost gratitude to the divine Almighty God for the boundless love and the bestowal of wisdom and knowledge that guided me in designing this report to completion.

I cannot forget to acknowledge my family, especially my husband, who has been my greatest support and source of encouragement throughout this journey to the completion of this research report.

Specifically, I want to thank my supervisor, Mrs. Atim Miria, for her indispensable guidance, which enabled me to successfully write this wonderful research report.

Appreciation to all the tutors of Jerusalem School of Nursing and Midwifery, my classmates, and friends.

I would also like to express my sincere gratitude to the Medical Superintendent of Nebbi hospital, the In-charge ANC, and all the staff of Nebbi hospital, especially those in

the maternity ward, for being there for me whenever I needed their assistance.

LIST OF ABBREVIATIONS

ANC: Antenatal Care

WHO: World Health Organization

MoH: Ministry of Health

RCT: Reproductive and Child Health

SPSS: Statistical Package for Social Sciences

HIV: Human Immunodeficiency Virus

EMTCT: Elimination of Mother-to-Child Transmission

MS: Medical Superintendent

AIDS: Acquired Immuno-deficiency Syndrome

SOURCE OF FUNDING

The study was not funded.

CONFLICT OF INTEREST

No conflict of interest has been declared.

AUTHOR BIOGRAPHY

Adero Teddy is a student of Nursing and Midwifery at Jerusalem Institute of Health Sciences.

Atim Miria is a research supervisor and a tutor of Midwifery at Jerusalem Institute of Health Sciences.

REFERENCES

- 1) Annie, C. (2021). Factors influencing male involvement in maternal child healthcare (mch) in solwezi district.
- 2) Auma, I., Nabaweesi, D., Orech, S., Alege, J. B., & Komakech, A. (2023). Determinants of male involvement in antenatal care at Palabek Refugee Settlement, Lamwo district, Northern Uganda. *BMC Pregnancy and Childbirth*, 23, 325. <https://doi.org/10.1186/s12884-023-05617-2>
- 3) Babughirana, G., Gerards, S., Mokori, A., Nangosha, E., Kremers, S., & Gubbels, J. (2020). Maternal and newborn healthcare practices: Assessment of the uptake of lifesaving services in Hoima District, Uganda. *BMC Pregnancy and Childbirth*, 20(1), 686. <https://doi.org/10.1186/s12884-020-03385-x>
- 4) Gibore, N. S., Bali, T. A. L., & Kibusi, S. M. (2019). Factors influencing men's involvement in antenatal care services: A cross-sectional study in a low resource setting, Central Tanzania. *Reproductive Health*, 16(1), 52. <https://doi.org/10.1186/s12978-019-0721-x>
- 5) Kabanga, E., Chibwae, A., Basinda, N., & Morona, D. (2019). Prevalence of male partners involvement in antenatal care visits – in Kyela district, Mbeya. *BMC Pregnancy and Childbirth*, 19(1), 321. <https://doi.org/10.1186/s12884-019-2475-4>
- 6) Kirui, N. J. (2021). Male participation in utilization of antenatal care services among their spouses in selected manufacturing industries in nairobi city county, kenya.
- 7) Mamo, Z. B., Kebede, S. S., Agidew, S. D., & Belay, M. M. (n.d.). Determinants of Male Partner Involvement during Antenatal Care among Pregnant Women in Gedeo Zone, South Ethiopia: A Case-Control Study. *Annals of Global Health*, 87(1), 19. <https://doi.org/10.5334/aogh.3003>
- 8) Muia, P. K., Mbuthia, G. W., & Mugambi, R. K. (2022a). Determinants of male partner involvement in antenatal care services at Kangundo Sub-County Hospital in Kenya. *African Health Sciences*, 22(3), 93–99. <https://doi.org/10.4314/ahs.v22i3.11>
- 9) Manda-Taylor, L., Mwale, D., Phiri, T., Walsh, A., Matthews, A., Brugha, R., ... & Byrne, E. (2017). Changing times? Gender roles and relationships in maternal, newborn and child health in Malawi. *BMC pregnancy and childbirth*, 17(1), 321.
- 10) WHO (2015). Trends in maternal Mortality:1990 to 2015 Estimates by WHO, UNICEF, UNFPA, World Bank Group and United Nations Population Division Executive summary.2015.(Google Scholar)
- 11) Yargawa, J., & Leonardi-Bee, J. (2015). Male involvement and maternal health outcomes: systematic review and meta-analysis. *J Epidemiol Community Health*, 69(6), 604-612.

12) Atuahene, M. D., Arde-acquah, S., Atuahene, N. F., Adjui, M., & Ganle, J. K. (2017). Inclusion of men in maternal and safe motherhood services in inner-city communities in Ghana: evidence from a descriptive cross-sectional survey, 1–10. HYPERLINK "https://doi.org/10.1186/s12884-017-1590-3" https://doi.org/10.1186/s12884-017-1590-3

13) Byamugisha, R., Åström, A. N., Ndezi, G., Karamagi, C. A. S., & Tylleskär, T. (2017). Male

partner antenatal attendance and HIV testing in eastern Uganda: A randomized facility-based intervention trial. Journal of the International AIDS Society, 13(1), 16.

14) Ditekemena et al (2016). Determinants of male involvement in maternal and child health services in sub-Saharan Africa: a review. Reproductive health.

PUBLISHER DETAILS

Burundi Publishing



Burundi Publishing

Contact: +257 6266 2725

Email: burundipublishing@gmail.com

Website: <https://burundipublishing.com>

**Address: Avenue de l'université, Quartier Rohero I,
Bujumbura, Burundi**